

Phytochemical Constituents of *Nelumbo nucifera*

Ki Hyun Kim^{1,§}, Sang Wook Chang^{1,§}, Shi Yong Ryu², Sang Un Choi², and Kang Ro Lee^{1,*}

¹Natural Products Laboratory, College of Pharmacy, Sungkyunkwan University, Suwon 440-746, Korea

²Korea Research Institute of Chemical Technology, Daejeon 305-600, Korea

Abstract – Phytochemical investigation of the MeOH extract of the leaves of *Nelumbo nucifera* resulted in the isolation of five norsesquiterpenes, four flavonoids, two triterpenes and one alkaloid. Their chemical structures were characterized by spectroscopic methods to be (*E*)-3-hydroxymegastigm-7-en-9-one (**1**), (3*S*,5*R*,6*S*,7*E*)-megastigma-7-ene-3,5,6,9-tetrol (**2**), dendranthemoside B (**3**), icariside B₂ (**4**), sedumoside F₁ (**5**), luteolin (**6**), quercetin 3-*O*-β-D-glucuronide (**7**), quercetin 3-*O*-β-D-glucoside (**8**), isorhamnetin 3-*O*-rutinoside (**9**), alphitolic acid (**10**), maslinic acid (**11**), and *N*-methylasimilobine (**12**). Norsesquiterpenoids (**1-5**) and triterpenes (**10-11**) were isolated for the first time from this plant. Compounds **6** and **10-12** exhibited considerable cytotoxicity against four human cancer cell lines *in vitro* using a SRB bioassay.

Keywords – *Nelumbo nucifera*, Nymphaeaceae, Norsesquiterpenes, Cytotoxicity

Introduction

Nelumbo nucifera GAERTN (Nymphaeaceae) is a perennial aquatic plant, which is distributed throughout Asia (Van Bergen *et al.*, 1997; Kim, 1996). The seeds of *N. nucifera* have been used in traditional medicine for the alleviation of fever and treatment of bleeding, dizziness and hematuria (Bensky and Gamble, 1993). Thus far, much works have been done on the phytochemical constituents of *N. nucifera*. Mainly, various alkaloids such as nuciferine, *N*-nornuciferine, roemerine, liensinine, neferine and (–)-1(*S*)-norcoclaurine, have been reported from this herb (Furukawa, 1996; Luo *et al.*, 2005; Kashiwada *et al.*, 2005; Wu *et al.*, 2004; Agnihotri *et al.*, 2008). Flavonoids and their glycosides including (+)-catechin, quercetin, kaempferol and nelumboside A have been also isolated (Kim *et al.*, 2001; Jung *et al.*, 2003; Hyun *et al.*, 2006). The extract of this herb exhibited anti-diabetic (Lee *et al.*, 2001), anti-oxidant (Hyun *et al.*, 2006) and anti-obesity activities (Ono *et al.*, 2006). As parts of our continuing search for bioactive constituents from Korean natural sources, we investigated the constituents of the leaves of *N. nucifera*. As a consequence, we isolated five norsesquiterpenes (**1-5**), four flavonoids (**6-9**), two triterpenes (**10** and **11**) and one alkaloid (**12**) from its MeOH extract. This paper is the first report of isolation of norsesquiterpenoids from

this plant source. The isolated triterpenes (**10-11**) were also isolated for the first time from this plant. All the isolated compounds were tested for their cytotoxic activities against four human cancer cell lines *in vitro* using a SRB bioassay.

Experimental

General – Melting points were determined on a Gallenkamp melting point apparatus and are uncorrected. Optical rotations were measured on a JASCO P-1020 Polarimeter. IR spectra were recorded on a Bruker IFS-66/S FT-IR spectrometer. UV spectra were recorded with a Shimadzu UV-1601 UV-Visible spectrophotometer. NMR spectra were recorded on a Varian UNITY INOVA 500 NMR spectrometer. EIMS and FABMS data were obtained on a JEOL JMS700 mass spectrometer, and LC-ESI/MS data on an Agilent 1100LC/MSD trap SL LC/MS. Preparative HPLC was performed using a Gilson 306 pump with a Shodex refractive index detector and Alltech Silica 5 μ column (250 × 22 mm) or Econosil® RP-18 10 μ column (250 × 22 mm). Silica gel 60 (Merck, 70~230 mesh and 230~400 mesh) was used for column chromatography. TLC was performed using Merck precoated Silica gel F₂₅₄ plates and RP-18 F_{254s} plates. The packing material for molecular sieve column chromatography was Sephadex LH-20 (Pharmacia Co.). Low-pressure liquid chromatography was performed over Merck LiChroprep Lobar®-A Si 60 (240 × 10 mm) or LiChroprep Lobar®-A RP-18 (240 × 10 mm)

*Author for correspondence

Tel: +82-31-290-7710; E-mail: krlee@skku.ac.kr

§ These authors contributed equally to this work.

columns with a FMI QSY-0 pump (ISCO). Sep-Pak[®] (Waters, Vac 6cc) and RediSep[®] (ISCO, C-18 Reverse Phase 4.3 g) were also used for column chromatography.

Plant materials – The leaves of *Nelumbo nucifera* (4.5 kg) were purchased from Kyungdong herbal market, Seoul, Korea, in July 2005. A voucher specimen (SKKU-2005-7) of the plant was deposited in the herbarium of the College of Pharmacy, Sungkyunkwan University, Suwon, Korea.

Extraction and isolation – The leaves of *N. nucifera* (4.5 kg) were extracted at room temperature with 80% MeOH and evaporated under reduced pressure to give a residue (656 g), which was dissolved in water (800 mL) and partitioned with solvent to give hexane (49 g), CH₂Cl₂ (16 g), EtOAc (10 g), and BuOH (30 g) soluble portions. The hexane fraction (49 g) was separated over a silica gel column (hexane : EtOAc = 7 : 1 – 1 : 1) to yield six fractions (H1 - H6). Fraction H5 (1.1 g) was further separated over an RP-C₁₈ silica gel column (60% MeOH) to yield ten fractions (H51 - H510). Fraction H53 (25 mg) was purified over a silica gel prep. HPLC (CH₂Cl₂ : MeOH = 90 : 1) to yield compound **1** (11 mg, *R_t* = 15.5 min). In turn, fraction H58 (25 mg) and H59 (42 mg) were purified over a silica gel prep. HPLC (hexane : EtOAc = 1 : 1) to give compounds **10** (10 mg, *R_t* = 17.0 min) and **11** (9 mg, *R_t* = 14.5 min), respectively. The EtOAc fraction (10 g) was separated over a silica gel column with a solvent system of CHCl₃ : MeOH (10 : 1 – 1 : 1) as the eluant to give sixteen fractions (E1 - E16). Fraction E10 (519 mg) was further separated over an RP-C₁₈ silica gel column (50% MeOH) and purified by repeating recrystallization using MeOH to give compound **6** (9 mg). In turn, fraction E16 (473 mg) was further separated over an RP-C₁₈ silica gel column (50% MeOH) and purified with RediSep[®] (50% MeOH) to afford compound **7** (6 mg). Similarly, the BuOH fraction (30 g) was separated over a silica gel column (CHCl₃ : MeOH : H₂O = 7 : 3 : 0.5) to give seven fractions (B1 - B7). Fraction B1 (1.0 g) was separated over a Sephadex LH-20 column (CH₂Cl₂ : MeOH = 1 : 1) and silica gel column (CHCl₃ : MeOH = 10 : 1), and purified with RediSep[®] (60% MeOH) to give compound **12** (7 mg). Fraction B3 was further separated over an RP-C₁₈ silica gel column (20% MeOH) to yield four fractions (B31 - B34). In turn, fraction B31 (178 mg) was purified with a silica Lobar A[®]-column (CH₂Cl₂ : MeOH = 8 : 1) and silica gel prep. HPLC (hexane : CHCl₃ : MeOH = 3 : 5 : 1) to give compound **2** (11 mg, *R_t* = 13.5 min). Fraction B33 (100 mg) was separated over an RP-C₁₈ silica Lobar A[®]-column (40% MeOH) and purified with a silica gel prep. HPLC (CH₂Cl₂ : MeOH = 8 : 1) to give compound **4** (6 mg, *R_t* =

13.0 min). Fraction B32 (230 mg) was separated over a silica Lobar A[®]-column (hexane : CHCl₃ : MeOH = 1 : 5 : 1) and purified with a silica gel prep. HPLC (EtOAc : MeOH : H₂O = 11 : 1 : 0.5) to give compound **3** (38 mg, *R_t* = 16.5 min). Fraction B34 (57 mg) was separated over RediSep[®] (80% MeOH) and purified with a silica gel prep. HPLC (CHCl₃ : MeOH = 11 : 1) to give compound **5** (5 mg, *R_t* = 12.5 min). Fraction B5 (1.8 g) was further separated over a silica gel column (EtOAc : MeOH : H₂O = 10 : 1 : 0.5) to give two fractions (B51 - B52). In turn, fraction B51 (20 mg) was purified with Sep-Pak[®] (CHCl₃ : MeOH = 5 : 1) to give compound **8** (5 mg). Fraction B52 (171 mg) was purified with an RP-C₁₈ prep. HPLC (45% MeOH) to give compound **9** (5 mg, *R_t* = 20.0 min).

(E)-3-Hydroxymegastigm-7-en-9-one (1) – Colorless gum; [α]_D²⁵: -7.7° (*c* 0.3, CHCl₃); FAB-MS *m/z* : 211 [M+H]⁺; ¹H-NMR (500 MHz, CDCl₃): δ 0.83 (3H, d, *J* = 6.5 Hz, H-13), 0.89 (3H, s, H-11), 0.92 (3H, s, H-12), 0.95 (1H, m, H-4a), 1.17 (1H, m, H-2a), 1.54 (1H, dd, *J* = 10, 11 Hz, H-6a), 1.67 (1H, m, H-5a), 1.79 (1H, m, H-2b), 2.08 (1H, m, H-4b), 2.27 (3H, s, H-10), 3.84 (1H, dddd, *J* = 4.5, 4.5, 11.5, 12.0 Hz, H-3), 6.07 (1H, d, *J* = 16.0 Hz, H-8), 6.53 (1H, dd, *J* = 10.0, 16.0 Hz, H-7); ¹³C-NMR (125 MHz, CDCl₃): δ 21.3 (C-13), 21.8 (C-11), 27.4 (C-10), 31.1 (C-5), 31.5 (C-12), 35.8 (C-1), 44.5 (C-4), 50.3 (C-2), 57.8 (C-6), 66.8 (C-3), 133.9 (C-8), 149.2 (C-7), 199.5 (C-9).

(3S,5R,6S,7E)-Megastigma-7-ene-3,5,6,9-tetrol (2) – Colorless gum; [α]_D²⁵: -22.7° (*c* 0.23, MeOH); FAB-MS *m/z* : 245 [M+H]⁺; ¹H-NMR (500 MHz, CD₃OD): δ 0.88 (3H, s, H-11), 1.11 (3H, s, H-13), 1.23 (3H, s, H-12), 1.27 (3H, d, *J* = 6.0 Hz, H-10), 1.47 (1H, m, H-2a), 1.68 (1H, t, *J* = 12.0 Hz, H-2b), 1.72 (1H, t, *J* = 12.0 Hz, H-4a), 1.75 (1H, m, H-4b), 4.05 (1H, m, H-3), 4.34 (1H, m, H-9), 5.79 (1H, dd, *J* = 6.0, 16.0 Hz, H-8), 6.09 (1H, dd, *J* = 1.0, 16.0 Hz, H-7); ¹³C-NMR (125 MHz, CD₃OD): δ 22.9 (C-10), 25.0 (C-12), 25.8 (C-13), 26.3 (C-11), 39.5 (C-1), 44.5 (C-4), 45.3 (C-2), 64.1 (C-3), 68.3 (C-9), 76.5 (C-5), 77.7 (C-6), 129.9 (C-7), 134.9 (C-8).

Dendranthemoside B (3) – Colorless gum; [α]_D²⁵: -42.1° (*c* 1.0, MeOH); FAB-MS *m/z* : 411 [M+Na]⁺; ¹H-NMR (500 MHz, CD₃OD): δ 0.83 (3H, d, *J* = 6.5 Hz, H-13), 0.89 (3H, s, H-11), 1.06 (3H, s, H-12), 1.51 (1H, m, H-4a), 1.60 (1H, m, H-2a), 1.69 (1H, m, H-2b), 1.87 (1H, m, H-4b), 2.12 (1H, m, H-5), 2.29 (3H, s, H-10), 3.14-3.88 (5H, m, sugar-H), 3.99 (1H, m, H-3), 4.38 (1H, d, *J* = 8.0 Hz, H-1'), 6.35 (1H, d, *J* = 16.0 Hz, H-8), 6.89 (1H, d, *J* = 16.0 Hz, H-7); ¹³C-NMR (125 MHz, CD₃OD): δ 15.3 (C-13), 23.9 (C-12), 24.7 (C-11), 26.2 (C-10), 34.1 (C-5), 36.6 (C-4), 39.7 (C-1), 41.2 (C-2), 61.6 (C-6'), 70.5 (C-

4'), 73.9 (C-2'), 74.2 (C-3), 76.6 (C-5'), 76.8 (C-3'), 77.8 (C-6), 101.5 (C-1'), 130.4 (C-8), 153.0 (C-7), 199.6 (C-9).

Icariside B₂ (4) – Colorless gum; $[\alpha]_D^{25}$: -69.8° (*c* 0.07, MeOH); FAB-MS *m/z* : 386 [M]⁺; ¹H-NMR (500 MHz, CD₃OD): δ 0.97 (3H, s, H-11), 1.17 (3H, s, H-12), 1.22 (3H, s, H-13), 1.43 (1H, m, H-2a), 1.74 (1H, m, H-4a), 1.80 (1H, m, H-2b), 2.30 (3H, s, H-10), 3.14-3.85 (5H, m, sugar-H), 3.87 (1H, m, H-3), 4.35 (1H, d, *J* = 8.0 Hz, H-1'), 6.20 (1H, d, *J* = 16.0 Hz, H-8), 7.17 (1H, d, *J* = 16.0 Hz, H-7); ¹³C-NMR (125 MHz, CD₃OD): δ 19.0 (C-13), 24.3 (C-12), 26.2 (C-10), 28.2 (C-11), 34.7 (C-1), 36.9 (C-4), 44.0 (C-2), 61.5 (C-6'), 67.1 (C-5), 69.9 (C-6), 70.4 (C-4'), 71.5 (C-3), 73.9 (C-2'), 76.7 (C-5'), 76.9 (C-3'), 101.7 (C-1'), 132.6 (C-8), 144.0 (C-7), 199.0 (C-9).

Sedumoside F₁ (5) – Colorless gum; $[\alpha]_D^{25}$: -3.3° (*c* 0.03, MeOH); FAB-MS *m/z* : 397 [M+Na]⁺; ¹H-NMR (500 MHz, CD₃OD): δ 0.82 (3H, d, *J* = 6.5 Hz, H-13), 0.88 (3H, s, H-11), 0.91 (3H, s, H-12), 0.92 (1H, m, H-4a), 1.11 (1H, m, H-2a), 1.28 (3H, d, *J* = 6.5 Hz, H-10), 1.32 (1H, m, H-6), 1.53 (1H, m, H-5), 1.69 (1H, m, H-2b), 1.96 (1H, H-4b), 3.14-3.80 (5H, m, sugar-H), 3.81 (1H, m, H-3), 4.35 (1H, d, *J* = 8.0 Hz, H-1'), 4.36 (1H, m, H-9), 5.35 (1H, dd, *J* = 10.0, 16.0 Hz, H-7), 5.53 (1H, dd, *J* = 7.0, 16.0 Hz, H-8); ¹³C-NMR (125 MHz, CD₃OD): δ 21.6 (C-10), 26.8 (C-11), 26.9 (C-13), 32.1 (C-5), 32.2 (C-12), 35.8 (C-1), 45.5 (C-4), 51.1 (C-2), 58.5 (C-6), 62.4 (C-6'), 67.3 (C-3), 71.2 (C-4'), 75.3 (C-2'), 77.9 (C-5'), 78.0 (C-3'), 78.0 (C-9), 102.2 (C-1'), 133.1 (C-7), 136.4 (C-8).

Luteolin (6) – Yellow amorphous powder; ESI-MS (negative) *m/z* : 285 [M-H]⁻; UV λ_{max} (MeOH) nm: 253, 267, 291 (sh), 349; IR ν_{max} (KBr): 3400, 1600, 1605, 1498 cm⁻¹; ¹H-NMR (500 MHz, CD₃OD): δ 6.22 (1H, s, H-6), 6.45 (1H, s, H-8), 6.54, (1H, s, H-3), 6.91 (1H, d, *J* = 8.5 Hz, H-5'), 7.37 (1H, br. s, H-2'), 7.38 (1H, br. s, H-6'); ¹³C-NMR (125 MHz, CD₃OD): δ 93.8 (C-8), 98.9 (C-6), 102.7 (C-3), 104.1 (C-10), 113.0 (C-6'), 115.6 (C-5'), 119.1 (C-2'), 122.5 (C-1'), 145.9 (C-3'), 149.8 (C-4'), 158.2 (C-9), 162.0 (C-5), 164.8 (C-7), 165.2 (C-2), 182.7 (C-4).

Quercetin 3-O- β -D-glucuronide (7) – Yellow amorphous powder; ESI-MS (negative) *m/z* : 477 [M-H]⁻; UV λ_{max} (MeOH) nm: 253, 267, 291; ¹H-NMR (500 MHz, CD₃OD): δ 3.31-3.95 (5H, m, sugar-H), 5.24 (1H, d, *J* = 8.0 Hz, H-1'), 6.22 (1H, s, H-6), 6.41, (1H, s, H-8), 6.85 (1H, d, *J* = 8.5 Hz, H-5'), 7.58 (1H, dd, *J* = 2.0, 8.5 Hz, H-6'), 7.60 (1H, d, *J* = 2.0 Hz, H-2'); ¹³C-NMR (125 MHz, CD₃OD): δ 71.4 (C-2''), 73.8 (C-3''), 75.6 (C-4''), 75.6 (C-5''), 93.6 (C-8), 98.8 (C-6), 101.4 (C-1''), 103.9 (C-10), 115.2 (C-2'), 116.2 (C-5'), 120.6 (C-1'), 120.9 (C-6'), 133.2 (C-3),

144.9 (C-3'), 148.6 (C-4'), 156.3 (C-2), 156.3 (C-9), 161.3 (C-5), 164.3 (C-7), 168.7 (C-6''), 177.2 (C-4).

Quercetin 3-O- β -D-glucoside (8) – Yellow amorphous powder; FAB-MS *m/z* : 465 [M+H]⁺; UV λ_{max} (MeOH) nm : 257, 264 (sh), 295, 355; IR ν_{max} (KBr): 3415, 1655, 1605, 1205 cm⁻¹; ¹H-NMR (500 MHz, CD₃OD): δ 5.25 (1H, d, *J* = 7.5 Hz, H-1''), 6.21 (1H, *J* = 2.0 Hz, H-6), 6.40 (1H, *J* = 2.0 Hz, H-8), 6.88 (1H, d, *J* = 8.0 Hz, H-5'), 7.59 (1H, dd, *J* = 8.0, 2.0 Hz, H-6'), 7.73 (1H, d, *J* = 2.0 Hz, H-2''); ¹³C-NMR (125 MHz, CD₃OD) : δ 61.3 (C-6''), 70.0 (C-4''), 74.6 (C-2''), 76.9 (C-3''), 77.2 (C-5''), 93.6 (C-8), 98.7 (C-6), 103.2 (C-1''), 104.5 (C-10), 114.9 (C-2'), 116.4 (C-5'), 121.7 (C-6'), 121.9 (C-1'), 134.5 (C-3), 144.6 (C-3'), 148.7 (C-4'), 157.3 (C-2), 157.8 (C-9), 161.8 (C-5), 164.9 (C-7), 178.3 (C-4).

Isorhamnetin 3-O-rutinoside (9) – Yellow amorphous powder; FAB-MS *m/z* : 625 [M+H]⁺; ¹H-NMR (500 MHz, CD₃OD): δ 1.10 (1H, d, *J* = 6.0 Hz, H-6''), 3.97 (1H, s, OCH₃), 4.53 (1H, s, H-1''), 5.24 (1H, d, *J* = 7.5 Hz, H-1''), 6.23 (1H, s, H-6), 6.43 (1H, s, H-8), 6.93 (1H, d, *J* = 8.0 Hz, H-5'), 7.65 (1H, d, *J* = 8.0 Hz, H-6'), 7.96 (1H, s, H-2''); ¹³C-NMR (125 MHz, CD₃OD): δ 16.7 (C-6''), 55.6 (C-3'-OCH₃), 67.3 (C-6''), 68.6 (C-5''), 70.4 (C-4''), 70.8 (C-2''), 70.9 (C-3''), 72.7 (C-4''), 74.7 (C-2''), 76.2 (C-5''), 77.0 (C-3''), 93.8 (C-8), 98.9 (C-6), 101.3 (C-1''), 103.0 (C-1''), 104.5 (C-10), 113.4 (C-2'), 114.9 (C-5'), 121.8 (C-6'), 122.7 (C-1'), 134.3 (C-3), 147.2 (C-3'), 149.6 (C-4'), 157.4 (C-2), 157.7 (C-9), 161.9 (C-5), 165.1 (C-7).

Alphitolic acid (10) – White powder, mp 275 - 278 °C; $[\alpha]_D^{25}$: $+176.8^\circ$ (*c* 0.065, CHCl₃); ESIMS *m/z* : 471 [M-H]⁻; ¹H-NMR (500 MHz, CDCl₃): δ 0.82 (3H, s, H-24), 0.92 (3H, s, H-25), 0.96 (3H, s, H-26), 0.99 (3H, s, H-23), 1.03 (3H, s, H-27), 1.71 (3H, s, H-30), 3.01 (1H, m, H-19), 3.70 (1H, m, H-2), 4.63 (1H, d, *J* = 1.5 Hz, H-29 β), 4.75 (1H, d, *J* = 1.5 Hz, H-29 α); ¹³C-NMR (125 MHz, CDCl₃): δ 14.9 (C-27), 16.3 (C-26), 16.7 (C-24), 17.6 (C-25), 18.5 (C-6), 19.6 (C-30), 21.2 (C-11), 25.6 (C-12), 28.7 (C-23), 29.9 (C-21), 30.8 (C-15), 32.4 (C-16), 34.4 (C-7), 37.2 (C-22), 38.5 (C-13), 38.8 (C-10), 39.4 (C-4), 41.0 (C-8), 42.7 (C-14), 47.0 (C-19), 47.1 (C-1), 49.5 (C-18), 50.7 (C-9), 55.7 (C-5), 56.5 (C-17), 69.8 (C-2), 84.1 (C-3), 110.3 (C-29), 150.5 (C-20), 177.0 (C-28).

Maslinic acid (11) – White powder, mp 241 - 245 °C; $[\alpha]_D^{25}$: $+45.1^\circ$ (*c* 0.17, MeOH); IR λ_{max} (CHCl₃) : 3400, 2910, 1675 cm⁻¹; EIMS *m/z* : 472 (M⁺), 426, 408, 248, 223, 203; ¹H-NMR (500 MHz, CD₃OD): δ 0.81 (3H, s, H-29), 0.82 (3H, s, H-26), 0.91 (3H, s, H-25), 0.95 (3H, s, H-30), 1.01 (3H, s, H-24), 1.02 (3H, s, H-27), 1.17 (3H, s, H-23), 2.91 (1H, d, *J* = 10.0 Hz, H-3), 3.62 (1H, m, H-2), 5.27 (1H, t, *J* = 3.5 Hz, H-12); ¹³C-NMR (125 MHz, CD₃OD):




Fig. 1. The structures of 1-12 from *N. nucifera*

δ 16.4 (C-25), 17.1 (C-24), 17.2 (C-26), 18.4 (C-6), 23.3 (C-30), 23.3 (C-16), 23.5 (C-11), 25.7 (C-27), 27.8 (C-15), 28.9 (C-23), 30.5 (C-20), 32.7 (C-22), 32.8 (C-29), 32.8 (C-7), 33.8 (C-21), 38.0 (C-10), 39.3 (C-8), 39.4 (C-4), 41.5 (C-18), 41.7 (C-14), 46.0 (C-19), 46.2 (C-17), 47.3 (C-1), 47.7 (C-9), 55.4 (C-5), 68.0 (C-2), 83.2 (C-27), 121.7 (C-12), 144.4 (C-13), 179.0 (C-30).

N-Methylasimilobine (12) – Brown gum; FAB-MS m/z : 282 $[M+H]^+$; $[\alpha]_D^{25}$: -211.2° (c 0.66, $CHCl_3$); UV λ_{max} (MeOH) nm: 253, 267, 291 (sh), 349; IR λ_{max} (KBr): 3400, 1600, 1589, 1570, 1496, 1471, 1452, 1423, 1373, 1298, 1273, 1242, 1173, 1142 cm^{-1} ; 1H -NMR (500 MHz,

CD_3OD): δ 2.53 (3H, s, N- CH_3), 3.01 (2H, m, H-4), 3.12 (2H, m, H-5), 3.18 (2H, m, H-7), 3.56 (3H, s, 1-O CH_3), 6.64 (1H, s, H-3), 7.21 (1H, m, H-10), 7.24 (1H, m, H-9), 7.31 (1H, m, H-8), 8.30 (1H, d, J = 7.5 Hz, H-11); ^{13}C -NMR (125 MHz, CD_3OD): δ 28.0 (C-4), 34.4 (C-7), 42.6 (C-6, N- CH_3), 53.1 (C-5), 59.2 (C-1, O CH_3), 62.7 (C-6a), 115.0 (C-3), 126.0 (C-1a), 126.5 (C-1b), 127.0 (C-11), 127.4 (C-9), 127.6 (C-10), 127.9 (C-8), 128.9 (C-3a), 132.1 (C-11a), 135.9 (C-7a), 144.1 (C-1), 150.0 (C-2).

Test for cytotoxicity *in vitro* – Sulforhodamin B bioassay (SRB) was used as for cytotoxicity screening (Skehan *et al.*, 1990). The *in vitro* cytotoxicity of each

compound against four cultured human tumor cells was assessed at the Korean Research Institute of Chemical Technology. The cell lines used were A549 (non small cell lung adenocarcinoma), SK-OV-3 (ovarian cancer cells), SK-MEL-2 (skin melanoma), and HCT15 (colon cancer cells). Doxorubicin was used as a positive control. The cytotoxicities of doxorubicin against A549, SK-OV-3, SK-MEL-2, and HCT cell lines were IC_{50} 0.007, 0.056, 0.117, and 0.164 μ M, respectively.

Results and Discussion

Compounds **2-4**, and **6-12** were identified by comparing the 1H -, ^{13}C -NMR, and MS spectral data with the literature values to be (3*S*,5*R*,6*S*,7*E*)-megastigma-7-ene-3,5,6,9-tetrol (**2**) (Kijima *et al.*, 1996), dendranthemside B (**3**) (Pauli *et al.*, 1990), icariside B₂ (**4**) (Otsuka *et al.*, 1995), luteolin (**6**) (Kwak *et al.*, 2009), quercetin 3-*O*- β -D-glucuronide (**7**) (Ishimatsu *et al.*, 1989), quercetin 3-*O*- β -D-glucoside (**8**) (Kim, 2001), isorhamnetin 3-*O*-rutinoside (**9**) (Xu *et al.*, 2006), aliphatic acid (**10**) (Bae *et al.*, 1996), maslinic acid (**11**) (Kim *et al.*, 2005), and *N*-methylasimilobine (**12**) (Han *et al.*, 1989). The terpenoids **1-5**, and **10-11** were isolated for the first time from this plant. The following describes the structural elucidation of compounds **1** and **5**, which were for the second time isolated from natural sources.

Compound **1** was obtained as a colorless gum with a negative optical rotation ($[\alpha]_D^{25}$: -7.7° in $CHCl_3$). From the FAB-MS (m/z 211 $[M+H]^+$) and 1H - and ^{13}C -NMR spectral data, the molecular formula of **1** was deduced to be $C_{13}H_{22}O_2$. The 1H -NMR spectrum showed four methyls at δ 0.83 (3H, d, $J = 6.5$ Hz, H-13), 0.89 (3H, s, H-11), 0.92 (3H, s, H-12), and 2.27 (3H, s, H-10), one methine bearing an oxygen function at δ 3.84 (1H, dddd, $J = 4.5, 4.5, 11.5, 12.0$ Hz, H-3), and an *trans*-olefinic protons at δ 6.07 (1H, d, $J = 16.0$ Hz, H-8), and 6.53 (1H, dd, $J = 10.0, 16.0$ Hz, H-7). The ^{13}C -NMR spectrum demonstrated the presence of 13 carbon signals, consisting of one carbonyl carbon signal at δ 199.5, and two olefinic carbon signals at δ 133.9 and 149.2, which indicated the presence of an α,β -unsaturated ketone moiety. These spectral data suggested that **1** was an ionane-type norsesquiterpene (Weyerstahl *et al.*, 1994). The coupling pattern (dddd, $J = 4.5, 4.5, 11.5, 12.0$ Hz) of H-3 implied that the configuration of OH group at C-3 was an equatorial form based on comparison with published data (Weyerstahl *et al.*, 1994). This was also confirmed by the NOESY experiment, in which correlations of H-3 with H-5 and H-11, and of H-12 with H-13 were observed. Consequently, the structure of **1** was

Table 1. Cytotoxic activities of compounds (**1-12**) isolated from *N. nucifera*

Compound	IC_{50} (μ M) ^a			
	A549	SK-OV-3	SK-MEL-2	HCT-15
1	>30.0	>30.0	>30.0	>30.0
2	>30.0	>30.0	>30.0	>30.0
3	>30.0	>30.0	>30.0	>30.0
4	>30.0	>30.0	>30.0	>30.0
5	>30.0	>30.0	>30.0	>30.0
6	12.1	9.8	7.0	9.7
7	>30.0	>30.0	>30.0	>30.0
8	>30.0	>30.0	>30.0	>30.0
9	>30.0	>30.0	>30.0	>30.0
10	10.0	13.5	12.1	8.1
11	13.7	16.5	15.8	11.7
12	13.1	14.6	6.0	2.8
Doxorubicin	0.007	0.056	0.117	0.164

^a IC_{50} value of compounds against each cancer cell line, which was defined as the concentration (μ M) that caused 50% inhibition of cell growth *in vitro*.

determined to be (*E*)-3-hydroxymegastigm-7-en-9-one.

Compound **5** was obtained as a colorless gum with a negative optical rotation ($[\alpha]_D^{25}$: -3.3° in MeOH). From the FAB-MS (m/z 397 $[M+Na]^+$) and 1H - and ^{13}C -NMR spectral data, the molecular formula of **5** was deduced to be $C_{19}H_{34}O_7$. The 1H -NMR spectrum showed four methyls at δ 0.82 (3H, d, $J = 6.5$ Hz, H-13), 0.88 (3H, s, H-11), 0.91 (3H, s, H-12), and 1.28 (3H, d, $J = 6.5$ Hz, H-10), two methines bearing an oxygen function at δ 3.81 (1H, m, H-3), and 4.36 (1H, m, H-9), an *trans*-olefinic protons at δ 5.35 (1H, dd, $J = 10.0, 16.0$ Hz, H-7), and 5.53 (1H, dd, $J = 7.0, 16.0$ Hz, H-8), together with β -D-glucopyranosyl moiety at 3.14-3.80 (5H, m, sugar-H), and 4.35 (1H, d, $J = 8.0$ Hz, H-1'). The ^{13}C -NMR spectrum demonstrated the presence of 19 carbon signals, consisting of two olefinic carbon signals at δ 133.1 and 136.4, and two oxygenated carbon signals at δ 67.3 and 78.0. An anomeric carbon signal at δ 102.2 and five oxygenated carbon signals (δ 78.0, 77.9, 75.3, 71.2, and 62.4) suggested the presence of glucopyranoside (Stephen *et al.*, 1977). The coupling constant ($J = 8.0$ Hz) of the anomeric proton of D-glucose indicated that it was β -form (Stephen *et al.*, 1977). The HMBC correlation between H-1' (δ 4.35) and C-9 (δ 78.0) indicated that sugar moiety was located at C-9. In the NOESY experiment, NOESY correlations between H-3 and H-5, H-11 and between H-11 and H-5, H-7 and between H-12 and H-13 were observed. On the basis of above information and optical rotation ($[\alpha]_D^{25}$: -3.3°) of **5** in comparison with published data ($[\alpha]_D$: -11.2°) (Morikawa

et al., 2007), the stereochemistry of carbon at C-3, 5, 6, and 9 was assigned to be *S*, *R*, *S*, and *R* form, respectively. Based on further comparison with published data (Morikawa *et al.*, 2007), the structure of **5** was identified as sedumoside F₁.

The isolated compounds (**1** - **12**) were tested *in vitro* for cytotoxicity against the A549, SK-OV-3, SK-MEL-2, and HCT15 human tumor cells using the SRB assay. Compounds **6** and **10-12** exhibited considerable cytotoxicity against the A549, SK-OV-3, SK-MEL-2, and HCT15 cell lines as shown in Table 1.

Acknowledgements

The authors would like to thank Mr. Do Kyun Kim, Dr. Eun Jung Bang, and Dr. Jung Ju Seo at the Korea Basic Science Institute for the NMR and MS spectra measurements.

References

- Agnihotri, V.K., ElSohly, H.N., Khan, S.I., Jacob, M.R., Joshi, V.C., Smillie, T., Khan, I.A., and Walker, L.A., Constituents of *Nelumbo nucifera* leaves and their antimalarial and antifungal activity. *Phytochemistry Letters* **1**, 89-93 (2008).
- Bae, K.H., Lee, S.M., Lee, E.S., Lee, J.S., and Kang, J.S., Isolation and quantitative analysis of betulinic acid and aliphatic acid from *Zyziphi Fructus*. *Yakhak Hoeji* **40**, 558-562 (1996).
- Bensky, D. and Gamble, A., *Chinese Herbal Medicine*, Materia Medica revised edition, Eastland press, p. 262 (1993).
- Furukawa, H., Studies on the alkaloids of *Nelumbo nucifera* GAERTN. NMR spectra of liensinine type alkaloids. *Yakugaku Zasshi* **86**, 883-886 (1996).
- Han, B.H., Park, M.H., and Han, Y.N., Aporphine and tetrahydrobenzylisoquinoline alkaloids from the seeds of *Zizyphus vulgaris* var. *spinosa*. *Arch. Pharm. Res.* **12**, 263-268 (1989).
- Hyun, S.K., Jung, Y.J., Chung, H.Y., Jung, H.A., and Choi, J.S., Isohammetin glycosides with free radical and ONOO⁻ scavenging activities from the stamens of *Nelumbo nucifera*. *Arch. Pharm. Res.* **29**, 287-292 (2006).
- Ishimatsu, M., Tanaka, T., Nonaka, G.I., and Nishioka, I., Alnusins A and B from the leaves of *Alnus sieboldiana*. *Phytochemistry* **28**, 3179-3184 (1989).
- Jijima, H., Otsuka, H., Ide, T., Ogimi, C., Hirata, E., Takushi, A., and Takeda, Y., Glycosides of megastigmane and of the simple alcohols from *Alangium premnifolium*. *Phytochemistry* **42**, 723-727 (1996).
- Jung, H.A., Kim, J.E., Chung, H.Y., and Choi, J.S., Antioxidant principles of *Nelumbo nucifera* stamens. *Arch. Pharm. Res.* **26**, 279-285 (2003).
- Kashiwada, Y., Aoshima, A., Ikeshiro, Y., Chen, Y.P., Furukawa, H., Itoigawa, M., Fujioka, T., Mihashi, K., Cosentino L, M., Morris-Natschke, S.L., and Lee, K.H., Anti-HIV benzylisoquinoline alkaloids and flavonoids from the leaves of *Nelumbo nucifera*, and structure-activity correlations with related alkaloids. *Bioorg. Med. Chem.* **13**, 443-448 (2005).
- Kim, J.S., Cho, S.M., Kim, J.H., and Lee, M.W., Phenolic compounds from the node of lotus rhizome (*Nelumbo nucifera* Gaertn). *Yakhak Hoechi* **45**, 599-603 (2001).
- Kim, D.K., Antioxidant components from the aerial parts of *Lactuca scariola* L. *Arch. Pharm. Res.* **24**, 427-430 (2001).
- Kim, D.H., Han, K.M., Chung, I.S., Kim, D.K., Kim, S.H., Kwon, B.M., Jeong, T.S., Park, M.H., Ahn, E.M., and Baek, N.I., Triterpenoids from the flower of *Campsis grandiflora* K. Schum. as human Acyl-CoA: cholesterol acyltransferase inhibitors. *Arch. Pharm. Res.* **28**, 550-556 (2005).
- Kim, T.J., *Korean Resources Plants*. Seoul Natural University Publisher, Seoul, Korea, Vol. II, p.95 (1996).
- Kwak, J.H., Kim, H.J., Lee, K.H., Kang, S.C., and Zee, O.P., Antioxidative iridoid glycosides and phenolic compounds from *Veronica peregrina*. *Arch. Pharm. Res.* **32**, 207-213 (2009).
- Lee, M.W., Kim, J.S., Cho, S.M., Kim, J.H., and Lee, J.S., Anti-diabetic constituent from the node of lotus rhizome (*Nelumbo nucifera* Gaertn). *Nat. Prod. Sci.* **7**, 107-109 (2001).
- Luo, X., Chen, B., Liu, J., and Yao, S., Simultaneous analysis of *N*-normuciferine, *O*-normuciferine, nuciferine, and roemerine in leaves of *Nelumbo nucifera* Gaertn by high-performance liquid chromatography-photodiode array detection-electrospray mass spectrometry. *Anal. Chim. Acta* **538**, 129-133 (2005).
- Morikawa, T., Zhang, Y., Nakamura, S., Matsuda, H., Muraoka, O., and Yoshikawa, M., Bioactive constituents from Chinese natural medicines. XXII.¹⁾ Absolute structures of new megastigmane glycoside, sedumosides E₁, E₂, F₁, F₂, and G, from *Sedum sarmentosum* (Crassulaceae). *Chem. Pharm. Bull.* **55**, 435-441 (2007).
- Ono, Y., Hattori, E., Fukaya, Y., Imai, S., and Ohizumi, Y., Anti-obesity effect of *Nelumbo nucifera* leaves extract in mice and rats. *J. Ethnopharmacol.* **106**, 238-244 (2006).
- Otsuka, H., Kamada, K., Yao, M., Yuasa, K., Kida, I., and Takeda, Y., Alangionosides C-F, megastigmane glycosides from *Alangium premnifolium*. *Phytochemistry* **38**, 1431-1435 (1995).
- Pauli, N., Sequin, U., and Walter, A., Boscialin and boscialin 4'-*O*-glucoside, two new compounds isolated from the leaves of *Boschia salicifolia* OLIV. *Helv. Chim. Acta* **73**, 578-582 (1990).
- Skehan, P., Storereng, R., Scudiero, D., Monks, A., McMahon, J., Vistica, D., Warren, J. T., Bokesch, H., Kenney, S., and Boyd, M.R., New colorimetric cytotoxicity assay for anticancer-drug screening. *J. Natl. Cancer Inst.* **82**, 1107-1112 (1990).
- Stephen, J.P., Louise, N.J., and David, C.P., High-resolution ¹H- and ¹³C-NMR Spectra of D-glucopyranose, 2-acetamido-2-deoxy-D-glucopyranose, and related compounds in aqueous media. *Carbohydr. Res.* **59**, 19-34 (1977).
- Van Bergen, P.F., Hatcher, P.G., Boon, J.J., Collinson, M.E., and De Leeuw, J.W., Macromolecular composition of the propagule wall of *Nelumbo nucifera*. *Phytochemistry* **45**, 601-617 (1997).
- Weyerstahl, P., Marschall, H., Bork, W.R., and Rilk, R., Megastigmanes and other constituents of the absolute of *Boronia megastigma* from tasmania. *Liebigs Ann. Chem.* 1043-1047 (1994).
- Wu, S., Sun, C., Cao, X., Zhou, H., Hong, Z., and Pan, Y., Preparative counter-current chromatography isolation of liensinine and its analogues from embryo of the seed of *Nelumbo nucifera* GAERTN. using upright coil planet centrifuge with four multilayer coils connected in series. *J. Chromatogr. A* **1041**, 153-162 (2004).
- Xu, M.L., Moon, D.C., Lee, C.S., Woo, M.H., Lee, E.S., Jahng, Y., Chang, H.W., Lee, S.H., and Son, J.K., Cytotoxicity and DNA topoisomerase inhibitory activity of constituents isolated from the fruits of *Evodia officinalis*. *Arch. Pharm. Res.* **29**, 541-547 (2006).

Received May 8, 2009

Revised June 13, 2009

Accepted June 15, 2009